



## MEETING ABSTRACT

## Open Access

# Influence of INF-gamma gene polymorphism in HTLV-1 proviral load

Rodrigo Haddad<sup>1,2\*</sup>, Maurício C Rocha-Junior<sup>1,3</sup>, Daiani C Cilião-Alves<sup>2</sup>, Virgínia M D Wagatsuma<sup>1</sup>, Oswaldo M Takayanagui<sup>2</sup>, Eduardo A Donadi<sup>2</sup>, Dimas T Covas<sup>1,2</sup>, Simone Kashima<sup>1,3</sup>

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses  
Leuven and Gembloux, Belgium. 5-8 June 2011

## Background

Interferon-gamma (INF- $\gamma$ ) is a key cytokine involved in the defense against intracellular pathogens, such as HTLV-1, by coordinating the expression of immunologically relevant genes. Previous studies showed the importance of this cytokine on HTLV-1-related pathogenesis. Thus, we have investigated the possible association between INF- $\gamma$  gene single-nucleotide polymorphism linked to high and low producer phenotypes (INF- $\gamma$  [+874T(high)  $\rightarrow$  A(low)]) and risk of development of symptoms related with HTLV-1 infection and proviral load.

## Methods

The polymorphism +874 T/A of INF- $\gamma$  was analyzed by PCR-SSP in 93 patients HTLV-1 positive (HAC + HAM), stratified according to the presence (HAM, n = 50) or not (HAC, n = 43) of symptoms, and healthy controls (n = 150). Proviral load of infected patients (HAC and HAM) was determined by real-time PCR.

## Results

No significant difference was observed for allelic and genotypic frequencies of the +874T/A polymorphism of INF- $\gamma$  when correlated with HAC, HAM and healthy controls groups. The median of proviral load was lower in HAC than HAM group (p=0.0131). Also, the p-value is very close to significance (p=0.0523) for +874TT genotype (high producer of INF- $\gamma$ ) and low proviral load, compared to the genotypes +874AA and +874AT.

## Conclusion

Despite the lack of significant associations, the low proviral load appears correlated with high producer of INF- $\gamma$  (+874TT) genotype. Increasing the number of patients may lead to statistical relationship of the +874TT genotype with low proviral load.

## Acknowledgements

Financial Support FAPESP, CNPq, CTC/FUNDHERP and INCTC.

## Author details

<sup>1</sup>Regional Blood Center of Ribeirão Preto, Ribeirão Preto, Brazil. <sup>2</sup>Faculty of Medicine of Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil.

<sup>3</sup>Faculty of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A119

**Cite this article as:** Haddad et al.: Influence of INF-gamma gene polymorphism in HTLV-1 proviral load. *Retrovirology* 2011 **8**(Suppl 1):A119.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [rodrigohaddad@hemocentro.fmrp.usp.br](mailto:rodrigohaddad@hemocentro.fmrp.usp.br)

<sup>1</sup>Regional Blood Center of Ribeirão Preto, Ribeirão Preto, Brazil

Full list of author information is available at the end of the article